

Scotland's Rural College

## Variation in light interception traits in European spring barley landraces

Florence, A; Ennos, RA; Hoad, SP; Hoebe, PN

*Published in:*  
Field Crops Research

*DOI:*  
[10.1016/j.fcr.2019.06.006](https://doi.org/10.1016/j.fcr.2019.06.006)

Print publication: 01/09/2019

*Document Version*  
Peer reviewed version

[Link to publication](#)

*Citation for pulished version (APA):*  
Florence, A., Ennos, RA., Hoad, SP., & Hoebe, PN. (2019). Variation in light interception traits in European spring barley landraces. *Field Crops Research*, 241, [107549]. <https://doi.org/10.1016/j.fcr.2019.06.006>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1   **Title:**

2   **Variation in light interception traits in European spring barley landraces**

3   **Authors:**

4   **Anna Florence<sup>\*a</sup>, Richard A. Ennos<sup>b</sup>, Steve P. Hoad<sup>a</sup>, Peter N. Høebe<sup>a</sup>**

5   a Crop and Soil Systems Group, SRUC, West Mains Road, Edinburgh, EH9 3JG, UK

6   b Institute of Evolutionary Biology, School of Biological Sciences, Ashworth Laboratories, University of  
7   Edinburgh, Edinburgh EH9 3JT, UK

8   \*- corresponding author

9   **Corresponding author:**

10   Anna Florence ([Anna.Florence@sruc.ac.uk](mailto:Anna.Florence@sruc.ac.uk) )

11   **Funding Information:**

12   This work was supported by funding from the Scottish Government's Rural and Environmental Science  
13   and Analytical Services (RESAS) division, with support to Anna Florence through a PhD studentship.

14   **Declaration:**

15   Authors have no competing interests to declare

16

## **Abstract**

Improving the efficiency of photosynthesis is a potential strategy for increasing crop yields in the future, but this is only possible if genetic variation exists for this attribute within crop germplasm resources. A key component of photosynthetic efficiency is the plant's ability to intercept light. This study examined the extent of genetic variation, available within barley landraces from Europe, for parameters affecting light interception. Landraces varied in time spent between emergence and full canopy establishment, with those from Northern latitudes reaching canopy closure between 2 and 8 days faster than those from Southern latitudes. There was significant variation in leaf chlorophyll content between the landraces, but this was unrelated to site of origin. Landraces originating from locations with cooler temperature over the growing season held their leaves in a more planophile manner than those from warmer climates, resulting in a negative relationship between leaf angle and mean temperature at site of origin. We conclude that substantial genetic variation in key parameters affecting light interception have evolved among barley landraces in Europe that could be utilised in future breeding programmes to improve the efficiency of photosynthesis and increase crop yields.

## **Keywords:**

Photosynthesis; Landraces; Light Interception, Canopy Structure, Local Adaptation

## 1. Introduction

Cereal yield increases over the past century have mainly come about from improved harvest index (HI), fertiliser responsiveness and increased arable land area (Evans, 1997; Fischer and Edmeades, 2010; Reynolds et al., 2011, 2009). Yields are now stagnating in many areas of the world in staple crops such as wheat, maize and rice (Mackay et al., 2011; Ray et al., 2012). With potential arable area reaching a limit due to increasing pressure for land use, new avenues for increasing yield must come from increasing production per unit area of ground (Long et al., 2015; Zhu et al., 2010). Most modern cereal breeding programs are derived from a small number of parent plants. Within these breeding programmes there may not be sufficient genetic variation present to exploit novel traits for increasing yield. Breeders may therefore need to look more widely to identify sources of suitable variation.

This situation is well illustrated by barley (*Hordeum vulgare* L.), one of the founding crops of modern agriculture, with major uses in Europe including malt and animal feed. It is widely grown in Scotland with 1.39million tonnes of spring barley and 268,124 tonnes of winter barley produced in 2018 (The Scottish Government, 2018). Barley landraces represent a possible source of genetic variation that could be used for improving traits related to yield (Rodriguez et al., 2008; Villa et al., 2005). A landrace is defined as a 'heterogeneous (genetically and phenotypically variable) variety that is reproduced by farmers as populations that are subject to both artificial and natural selection' (Bellucci et al., 2013). In some marginal areas landraces have been seen to outperform conventional cultivars (Dwivedi et al., 2016; Yahiaoui et al., 2014) as they can be locally adapted to climatic conditions (Bellucci et al., 2013). Landraces have already been used successfully to introduce traits into maize and rice that have increased yield under drought and submergence conditions respectively (Bailey-Serres et al., 2010; Meseka et al., 2015, 2013; Xu and Mackill, 1996).

Yield formation can be summarised in an equation first proposed by Monteith (Monteith and Moss, 1977):

$$Y = 0.487 \cdot S_t \cdot \varepsilon_i \cdot \varepsilon_c \cdot \varepsilon_p$$

Where Y is yield,  $S_t$  is the total amount of incident solar radiation with 0.487 being the fraction which is photosynthetically active,  $\varepsilon_i$  the efficiency of the plant in intercepting the fraction of photosynthetically active radiation,  $\varepsilon_c$  the efficiency of the photosynthetic processes converting light to energy and  $\varepsilon_p$  the proportion of energy produced which is partitioned into harvestable product. Whilst  $\varepsilon_p$  has largely been optimised there is still potential to improve  $\varepsilon_i$  and  $\varepsilon_c$  through optimisation of light interception and photosynthetic reactions (Farquhar et al., 2001; Raines, 2011). Rate of canopy development, amount and arrangement of chlorophyll and the leaf canopy architecture are all characters which can contribute towards  $\varepsilon_i$  which may be targets for optimisation.

The rate of canopy development is one of the major factors affecting  $\varepsilon_i$  (Flood et al., 2011; Long et al., 2006; Nunes-Nesi et al., 2016). Early and rapid canopy establishment can allow crops to take advantage of shorter growing seasons in Northern latitudes where the greatest amount of radiation is available in early spring (Murchie et al., 2009; Parry et al., 2011; Richards, 2000; Zhu et al., 2010). Extending the duration of canopy maintenance with slower loss of chlorophyll content during the grain filling phase in 'Stay-green' varieties is associated with an increase in grain weight in barley, maize and wheat and can lead to increased yields (Diaz et al., 2005; Emebiri, 2013; Parry et al., 2011; Zheng et al., 2009).

The chlorophyll arrangement throughout the canopy will also affect the  $\varepsilon_i$  as canopy with a more even distribution of chlorophyll throughout the leaf layers along with a greater total volume of chlorophyll may increase total light captured by reducing the number of leaves becoming saturated in the upper layers of the canopy (Ort et al., 2011; Yin and Struik, 2015). Saturation of upper leaf layers is a limiting factor in light interception as the electron transport systems fall at relatively low light levels (Björkman and Demmig, 1987). A horizontal leaf arrangement leads to saturation of the upper canopy whereas a canopy with an upper leaf angle of  $75^\circ$  from the horizontal can have double the efficiency of energy

capture of a horizontal canopy at midday (Long et al., 2006). Leaf size also affects light interception and there is a trade-off between leaf size and self-shading (Amanullah et al., 2007; Long et al., 2006).

The primary aim of this study was to assess a collection of European spring barley landraces for variation in traits associated with light interception efficiency including the timing of canopy development, chlorophyll content and arrangement, leaf canopy architecture and HI. The secondary aim of the study was to relate any variation found in light interception traits to environmental conditions from the locations which the landraces originated in order to understand the factors that may have led to their local adaptation. From these tests, results are put into context of how trait variation can be considered for improvements in plant breeding and resilience to climate change.

## **2. Methods**

### **2.1 Seed source and Experimental Design**

The field experiment was carried out at Scotland's Rural College's Boghall farm in Midlothian, Scotland, UK (55°52'26"N 3°12'26"W) in spring and summer of 2014 and 2015. The soil type at these sites is a sandy loam (Macmerrey Series). The farm is situated on the south-east slope of the Pentland hills at an elevation of 190m and the previous crop in the fields on both years was spring barley.

The barley landrace material was collected from gene banks (Table 1) prior to the start of this project and the landraces were specifically chosen to represent a wide latitudinal range across Europe which encompasses a spread of different climatic conditions and season lengths. The latitude and longitude of their original collection was used for collection of climatic data (Table 2). The landraces were a mixture of 2 and 6 row types dependent on the number of rows of seeds present on each ear. The modern cultivar Concerto was included to represent modern pedigree bred germplasm as a comparison to the landraces and was included as it was the main variety in Scotland during the experimental years.

The experimental design was a fully randomised, blocked design with three blocks in 2014 and four in 2015 and twelve plots per block (Plots measured 0.5m<sup>2</sup>). Four replicate plants per plot were used as technical replicates. Plots were sown on 09/04/14 and 23/04/15 and each plot was treated with 120kg/hectare of nitrogen by hand with 60kg applied on 26/04/14 and 25/04/15 for respective years and an additional 60kg two weeks after the first application. An herbicide treatment was applied when the plants reached GS23 (Harmony 70g/ha + Oxytril 0.5L/ha + High load micra.m 1.0L/ha).

## **2.2 Climate**

The climate in the location of original collection for each of the landraces was included as a possible factor influencing the canopy structure. Climate data were obtained from the national meteorological offices in each country of origin (Table 2). The area over which the weather data was collected varies between countries from local weather data to regional data depending on the scale of reporting. It was always taken as the closest reported point to the latitude and longitude of origin of the landraces. The climatic variables reported are the total rainfall (mm) for spring/summer, the total number of sunlight hours for spring/summer and the average daily temperature (°C) for spring/summer. The data are long-term averages with FRA1, FIN1, BRI1 and SPN1 (Table 2) being from 1981-2010. GER2, NOR1, NOR2, CZE1 and GER3 and from 1961-1991 and ITA1 (Table 2) is from 1971-2000.

## **2.3 Crop Measurements**

### **2.3.1 Canopy establishment**

The Growth Stage (GS) of the plants were recorded weekly throughout the growth season and was assessed using the HGCA (AHDB) growth stage guide which is based on the Zadoks 100 point growth scale (HGCA (The Scottish Executive), 2006; Zadoks et al., 1974). The plot was deemed to have reached a specific growth stage when at least 50% of the plants in the plot had reached that growth stage. Additional, in depth, assessment of four replicate plants per block of the canopy structure including leaf angle, length and chlorophyll content (by proxy with SPAD readings) were measured at GS24, GS39 and GS59. At GS24 the plant is made up of the main shoot and four tillers and is in the

establishment phase of its lifecycle. By GS39 stem extension is underway and the flag leaf is fully emerged meaning that canopy establishment is complete. At GS59 the ear has fully emerged from the boot and the plant has progressed from vegetative growth to reproductive growth.

### **2.3.2 Chlorophyll content, Distribution and Leaf Dimensions**

The leaf chlorophyll content was assessed by proxy using of a SPAD meter (Minolta Corp, Ramsay, NJ). SPAD readings were taken on the uppermost leaf excluding the flag leaf on a weekly basis midway along the length of the leaf blade. SPAD readings at GS39 and 59 are reported here.

Leaf area was measured by detaching the leaves where they meet the stem and immediately passing them through a leaf area meter (Li-3100 are meter, LiCor Inc., Lincoln, NE) which calculated leaf area in cm<sup>2</sup>. Leaves were passed through the meter three times and the readings averaged. The leaves that had been used for leaf area measures were then placed in individual paper bags and dried in an oven (Ecocell, MMM Medcenter, Munich, Germany) at 80°C for 48 hours. The leaves were then weighed using a precision balance (Kern PLJ, D-72336, Kern & Sohn Gontbl, Balingen, Germany) in grams. The specific leaf area (SLA) was calculated as leaf area divided by leaf dry weight. The leaf area, dry weight and SLA were all measured at GS39 on the uppermost leaf excluding the flag leaf.

### **2.3.3 Leaf canopy architecture**

Leaf angle was measured at GS39 and 59 in relation to the stem directly above it using a Helix Oxford protractor (Maped Helix, West Midlands, UK) to the nearest 5°. Care was taken to avoid bending the leaf away from the stem by minimising handling prior to this measure being taken.

### **2.3.4 Allocation of resources**

Harvest took place on 12/08/14 and 04/09/15. The ears on the shoots used for the earlier structural measurements were individually hand threshed and the grain number, row count and grain weight recorded and the 1000grain weight calculated. Grain weight was measured using a precision balance



(Kern PLJ, D-72336, Kern & Sohn Gontbl, Balingen, Germany). The ear and the straw were harvested to allow calculation of harvest index. The straw was dried in an oven (Ecocell, MMM Medcenter, Munich, Germany) at 80°C for 48 hours and weighed using a precision balance (Kern PLJ, D-72336, Kern & Sohn Gontbl, Balingen, Germany). Harvest index was then calculated by dividing the grain weight by the combined weight of the grain plus the straw plus the chaff. A total yield in tonnes per hectare was not calculated for the lines as the experimental design limited the amount of material that could be collected for each plot.

## **2.4 Statistical Analysis**

We used an Analysis of Variance (ANOVA) model to determine whether there was a significant amount of variation among the landraces in each trait of interest at a significance level of  $p=0.05$ . Year was included as a factor to see if differences between the two years of the trial were present and if there was an interaction between the year of the trial and variation between the landraces in the trait of interest. The results of the ANOVA are reported as the test statistic F-value to show the ratio of between to within group variability with the degrees of freedom as a subscript followed by the p-value. Effects of year of the trial are also reported if significant. Regression analysis of climate and latitude with each measure of canopy structure was carried out to see if there was a relationship between traits and local climatic conditions of each landrace line. A multiple regression of leaf angle with latitude and temperature was used to examine if both factors regressed significantly with leaf angle. The linear regression of leaf angle with temperature is reported below as this was the significant factor. Results of the regression are reported as the test statistic t-value to show if the slope of the regression is significantly different from zero with the upper degree of freedom as a subscript followed by the p-value which is taken as significant at a level of  $p=0.05$ . In the regression analysis year was included in a factor to see if the response differed between years and this is reported where significant. All figures report an average of all data between the years. Correlation analysis was carried out on allocation of resources factors to assess if relationships were present between the variables.

The correlation coefficient of significant relationships is given followed by the p-value. All statistical analysis was carried out using GenStat 16<sup>th</sup> Edition (VSN International Ltd, Hemel Hempstead, UK).

### **3. Results**

#### **3.1 Growing conditions in experiments**

In 2014 the average daily temperature at Boghall ranged from 7°C to 16°C. The average monthly rainfall was 107mm and the average monthly hours of sunlight was 119 hours. In 2015 the average daily temperature ranged from 5.5°C to 14.5°C. The average monthly rainfall was 130mm and the average monthly hours of sunlight was 119 hours.

#### **3.2 Canopy Establishment**

Landraces differ significantly in their development rate between GS24 and GS39 ( $F_{10,52}=9.36$ ,  $p<0.001$ ) (Figure 1) ranging from 12-20 days. Year was included as a factor in the analysis and there was a difference between the two years of the trial ( $F_{10,52}=19.78$ ,  $p<0.001$ ) but there was no interaction effect between the landraces and the year. GS24-GS39 is the stage in the plant leading up to full canopy establishment where GS24 consists of a plant with the main shoot and four tillers. It then moves through stem extension until it reached GS39 where the flag leaf is fully emerged. The length of time spent between these growth stages declines significantly in length in landraces from higher latitudes ( $t_{20}=34.5$ ,  $p<0.001$ ,  $R^2=0.65$ ) with a significant effect when year was included as a factor ( $p<0.001$ ) in the regression analysis. The modern cultivar Concerto spent longer than the landraces to reach GS24 and a similar amount of time in the canopy establishment stage between GS24 and GS39 to the Southern European landraces.

#### **3.3 Chlorophyll Content, Distribution and Leaf Dimensions**

There were significant differences in SPAD readings between the lines at all three growth stages: GS24 ( $F_{11,57}=6.97$ ,  $p<0.001$ ), GS39 ( $F_{11,57}=4.45$ ,  $p<0.001$ ) and GS59 ( $F_{11,57}=2.07$ ,  $p=0.037$ ) (Figure 2) with SPAD values ranging from 32.5-43.4, 35.2-45.4 and 41.1-48.7 respectively. When year was included as a

factor in the ANOVA analysis it was seen that there was a significant difference between the years at GS24 ( $F_{1,57}=35.04$ ,  $p<0.001$ ) and GS59 ( $F_{1,57}=5.22$ ,  $p=0.026$ ). There was no relationship between either climate or latitude and leaf chlorophyll content. Concerto had higher SPAD readings than the landraces at all growth stages.

The length of the second leaf showed significant differences between the landraces at both GS39 ( $F_{11,57}=6.38$ ,  $p<0.001$ ) and GS59 ( $F_{11,57}=8.17$ ,  $p<0.001$ ) (Table 3) with leaf length between 23.2-30.3 and 20.8-29.7cm respectively. When year of trial was included as a factor in the analysis significant differences were present at both GS39 ( $F_{1,57}=53.32$ ,  $p<0.001$ ) and GS59 ( $F_{1,57}=34.71$ ,  $p<0.001$ ). There were no significant differences between the landraces in SLA (Table 3).

### **3.4 Leaf Canopy Architecture**

Landraces differ significantly in their leaf angle at GS39 ( $F_{11,57}=10.48$ ,  $p<0.001$ ) (Table 3) where the final leaf of the canopy has fully emerged and GS59 ( $F_{11,57}=14.74$ ,  $p<0.001$ ) (Table 3) where the ear has fully emerged and the plant is switching from the vegetative to reproductive phase of its lifecycle (Table 3). The leaf angles from vertical range from 18-45 degrees and 31-84 degrees respectively. When year of trial was included as a factor in the analysis significant effects were seen at GS39 only ( $F_{1,57}=14.69$ ,  $p<0.001$ ). Leaf angle increases significantly with average temperature and Fig. 3 shows this relationship at GS59 (which is when the ear is fully emerged and the canopy size is at its maximum) ( $t_{20}=28.47$ ,  $p<0.001$ ,  $R^2=0.56$ ) (Table 3) at the location of origin. The same relationship is present at the earlier GS39 where the flag leaf is fully emerged ( $t_{20}=12.31$ ,  $p=0.002$ ,  $R^2=0.35$ ) (Table 3). Concerto fitted into the pattern of the landraces when temperature of the trial site was used as their origin location.

### **3.5 Allocation of resources**

The HI showed significant differences among the landraces in the 2 row lines ( $F_{7,37}=23.72$ ,  $p<0.001$ ). The 1000 grain weight showed a significant difference between the landraces in both the 2 ( $F_{7,37}=8.12$ ,

p<0.001) and 6 row lines ( $F_{3,17}=7.57$ ,  $p=0.002$ ). When year was included as a factor in the analysis there was a significant difference in 1000 grain weight between the years of the trial ( $F_{1,37}=4.99$ ,  $p=0.032$ ). The number of grains per ear showed a significant difference between the landraces in the 2 row lines ( $F_{7,37}=5.88$ ,  $p<0.001$ ) (Table 4). When year was included as a factor in the analysis there was a significant difference in number of grains per ear between the years of the trial ( $F_{1,37}=48.57$ ,  $p<0.001$ ). There was a significant positive correlation of number of grains per ear with 1000 grain weight with a correlation coefficient of  $r=0.934$ ,  $p=0.001$  (Figure 4).

#### **4. Discussion**

The data collected in this study will allow a picture of the variation present in traits associated with photosynthetic efficiency in spring barley landraces to be assessed. It will also allow the variation seen to be examined for local adaptation to environmental conditions and how this variation can be subsequently used in pre-breeding programs. It was found that differences in rate of canopy development, chlorophyll content and canopy leaf angle all varied significantly between the landraces. Canopy development rate and leaf angle both varied with climatic conditions at the location of origin with shorter duration at higher latitudes and more planophile leaves at lower temperatures, suggesting adaptation to local condition. The traits looked at in this study were assessed out with the local climatic conditions where they originated which suggests that the variation found is under strong genetic control and this could be very beneficial to breeding programs.

##### **4.1 Canopy Establishment**

The Scandinavian landraces progressed quickly through stem extension to canopy closure (Figure 1) which would be advantageous for light interception, biomass production and grain development during the early phase of a shorter growing season. There were differences in the time spent in this phase of development between the years of the trial showing that the environment has an effect on development rate but there were also differences between the landraces which showed genetic variation is present between the lines. This has been seen in barley and other cereals including wheat and oats (Goyne et al., 1993; Kemanian et al., 2004; Muurinen and Peltonen-Sainio, 2006; Peltonen-

Sainio, 1997). Landraces from Northern latitudes will likely contain the photoperiod non-responsive polymorphism in the *Ppd-H1* gene (Jones et al., 2011; Turner et al., 2005) allowing them to move into flowering irrespective of day-length. This polymorphism has been linked to leaf size caused by changes in duration of leaf growth (Digel et al., 2016). Reaching full canopy establishment quicker may also be an advantage in out-competing weeds, shading out possible (weed) competitors in organic systems or allowing reduced herbicide application under conventional management (Sim et al. 2007; Kruk et al. 2006). The modern cultivar Concerto spent more time reaching GS24 than the landraces but progressed from GS24-GS39 at the same rate as the landraces from Southern European latitudes reaching full canopy closure later than the landraces. The early development of the Scandinavian landraces may be a trait of interest in developing new varieties which are able to take advantage of early light in a short growing season.

#### **4.2 Chlorophyll Content, Distribution and Leaf Dimensions**

During leaf emergence and through canopy closure there were differences in chlorophyll content between the landraces (Figure 2). Variation has been observed in modern varieties in a study of Sardinian wheat, barley and triticale (Giunta et al., 2002) with the varieties showing high levels of variation in chlorophyll content caused by a strong genetic and weak environmental and G\*E components suggesting that chlorophyll content has not been driven in a particular direction as a side effect of breeding for other traits such as plant height. An environmental effect was seen in this study as differences in chlorophyll content were seen at some growth stages between the years of the trial. The modern cultivar Concerto was included in this study and had higher SPAD readings than the landraces present. Sufficient variation may already exist for altering the volume of chlorophyll in barley in the current pool of parents but having landraces as an alternative allows options for wider genetic material to be introduced to the breeding programs. Maintaining chlorophyll content for longer may be of benefit, as in rice it was seen that for each extra day of canopy maintenance there was an increase of 0.2 tonnes per hectare in yield (Akita, 1989). Compared to modern cultivars, wheat landraces have been seen to begin to senesce quicker once they have reached grain filling which

suggests a potential negative side effect in using landraces in breeding (Gaju et al., 2016). This was visually observed in this study although senescence was not measured directly. This is something pre-breeding programs would need to be taken into account when using landraces.

The distribution of chlorophyll will be affected by the size and shape of the leaves and there were differences between the landraces in regard to leaf length at both GS39 and 59 but not in SLA suggesting that lines with a larger leaf surface area are thinner and *vice versa* (Table 3). This is supported by research in barley which showed no differences in SLA between cultivars (Giunta et al., 2002) in contrast to their findings in wheat and triticale which showed variation in SLA. Unfortunately due to experimental constraints caused by the small size of the trial plots it was not possible to measure the leaf area index or light interception. This would have completed the picture of how leaf size and shape is affecting the light capture of the landraces throughout different phases of growth and is something to be explored in future work. Selection for seedling leaves with a larger surface area has occurred in wheat and it was accompanied by an increased early plant biomass and vigour (Zhang et al., 2015). If a similar approach could be applied in barley then light interception efficiency could be increased early in the growth cycle especially in Northern latitudes where the growth season is short (Mukula and Rantanen, 1837).

#### **4.3 Leaf Canopy Architecture**

Leaf angle had been associated with cereal yields and our study showed significant variation in leaf angle among barley landraces (Table 3). Early studies in Maize showed a yield increase of 40% with a 10° leaf inclination from vertical (Pendleton et al., 1968). High yielding rice varieties such as 'Takanari' have been reported to have higher photosynthetic rates per leaf than other varieties (Taylaran et al., 2011) and an erect leaf posture (Nan Su San et al., 2018) along with decreased levels of photo-inhibition (Horton et al., 1999; Kumagai et al., 2014). The optimal crop ideotype has previously been that of an overall erect canopy (Donald, 1968) but it is now suggested that decreasing leaf angle from the bottom leaf layer of the canopy to the top would be more efficient in maximising light interception

(Ku et al., 2010; Long et al., 2006; Zhu et al., 2010). Rice hybrids are being developed with 5°, 10° and 20° flag, 1<sup>st</sup> and 2<sup>nd</sup> leaves respectively (Peng et al., 2008). Canopies could be developed not only with variation in leaf angle but also with differential volumes of chlorophyll through leaf layers tailored to local environmental conditions (Ort et al., 2015).

The landraces from Southern latitudes were characterised by an erectophile leaf angle which has a negative relationship with latitude and temperature (Table 3, Figure 3). As latitude and temperature may be related a multiple linear regression was used to try to untangle if leaf angle was responding to one or both of the factors and this showed that leaf angle was responding to temperature. In other work an erectophile canopy structure has been seen to be beneficial in coping with heat stress and increasing water-use- and photosynthetic-efficiency through reduction in heat loads (Ryel et al., 1993; Valladares and Pugnaire, 1999; Werner et al., 2001) reducing excess light levels causing photosynthetic saturation at midday (Falster and Westoby, 2003). This suggests a degree of local adaptation to climatic temperature and light levels in canopy structure although more work would be needed to confirm this. The modern cultivar Concerto which has been developed for a climate midway in the range seen for the landraces fitted well with the regression seen in the landraces with temperature possibly indicating that this pattern has been retained in new breeding material.

#### **4.4 Allocation of resources**

Variation among landraces in yield components and resource partitioning was observed (Table 4) and studies have found relationships between numbers of grain and grain weight (Acreche and Slafer, 2006; Calderini and Reynolds, 2000) with wheat showing a reduction in average grain weight with increasing numbers of grain (Acreche and Slafer, 2006). There is uncertainty over whether competition between grains for resources reduces weight when there are more grains present (Borrás et al., 2004). In the barley landraces, an increase in grain weight with grain number in the 2-row lines (Figure 4) may be a consequence of lower tiller numbers and more resources allocated per ear. Landrace total yields were not obtained on an area basis due to constraints created by the small plot size. This would

have been informative in understanding how canopy structure traits affects final yield. However, as landraces would need to enter a pre-breeding program to introduce traits of interest into new varieties high yields could be maintained through careful trait selection. Source-sink limitations will need to be considered when improving traits associated with photosynthetic efficiency as yields have been shown to be sink limited with the number of grain per m<sup>2</sup> being the major contributor to yield as opposed to grain weight (Burnett et al., 2016; Lynch et al., 2017; Madani et al., 2010; Serrago et al., 2013). In order for greater photosynthetic efficiency to enhance yields sink strength must be increased with higher number of floret production, higher numbers of productive tillers and the capacity for larger grains (Reynolds et al., 2009).

## **5. Acknowledgements**

The authors would like to thank the technical and trials staff at SRUC for their assistance with this work.

## **6. References**

- Acreche, M.M., Slafer, G.A., 2006. Grain weight response to increases in number of grains in wheat in a Mediterranean area. *F. Crop. Res.* 98, 52–59. <https://doi.org/10.1016/j.fcr.2005.12.005>
- Akita, S., 1989. Improving yield potential in tropical rice, in: *Progress in Irrigated Rice Research*. IRRI, Los Banos, pp. 41–73.
- Amanullah, Hassan, M.J., Nawab, K., Ali, A., 2007. Response of specific leaf area (SLA), leaf area index (LAI) and leaf area ratio (LAR) of Maize (*Zea mays*, L.) to plant density, rate and timing of nitrogen application. *World Appl. Sci. J.* 2, 235–243.
- Bailey-Serres, J., Fukao, T., Ronald, P., Ismail, A., Heuer, S., Mackill, D., 2010. Submergence Tolerant Rice: SUB1's Journey from Landrace to Modern Cultivar. *Rice* 3, 138–147. <https://doi.org/10.1007/s12284-010-9048-5>
- Bellucci, E., Bitocchi, E., Rau, D., Nanni, L., Ferradini, N., Giardini, A., Rodriguez, M., Attene, G., Papa,



352 R., 2013. Population structure of barley landrace populations and gene-flow with modern  
353 varieties. PLoS One 8, e83891. <https://doi.org/10.1371/journal.pone.0083891>

354 Björkman, O., Demmig, B., 1987. Photon yield of O<sub>2</sub> evolution and chlorophyll fluorescence  
355 characteristics at 77 K among vascular plants of diverse origins. Planta 170, 489–504.  
356 <https://doi.org/10.1007/BF00402983>

357 Borrás, L., Slafer, G.A., Otegui, M.E., 2004. Seed dry weight response to source-sink manipulations in  
358 wheat, maize and soybean: A quantitative reappraisal. F. Crop. Res. 86, 131–146.  
359 <https://doi.org/10.1016/j.fcr.2003.08.002>

360 Burnett, A.C., Rogers, A., Rees, M., Osborne, C.P., 2016. Carbon source–sink limitations differ  
361 between two species with contrasting growth strategies. Plant. Cell Environ. 39, 2460–2472.  
362 <https://doi.org/10.1111/pce.12801>

363 Calderini, D.F., Reynolds, M.P., 2000. Changes in grain weight as a consequence of de-graining  
364 treatments at pre- and post-anthesis in synthetic hexaploid lines of wheat. Aust. J. Agric. Res.  
365 27, 187–191. <https://doi.org/10.17700/jai.2015.6.1>

366 Diaz, C., Purdy, S., Christ, A., Morot-Gaudry, J.-F., Wingler, A., Masclaux-Daubresse, C., 2005.  
367 Characterization of Markers to Determine the Extent and Variability of Leaf Senescence in  
368 Arabidopsis. A Metabolic Profiling Approach. Plant Physiol. 138, 898–908.  
369 <https://doi.org/10.1104/pp.105.060764>

370 Digel, B., Tavakol, E., Verderio, G., Tondelli, A., Xu, X., Cattivelli, L., Rossini, L., von Korff, M., 2016.  
371 Photoperiod-H1 (Ppd-H1) Controls Leaf Size. Plant Physiol. 172, 405–415.  
372 <https://doi.org/10.1104/pp.16.00977>

373 Donald, C.M., 1968. The breeding of crop ideotypes. Euphytica 17, 385–403.  
374 <https://doi.org/10.1007/BF00056241>

375 Dwivedi, S.L., Ceccarelli, S., Blair, M.W., Upadhyaya, H.D., Are, A.K., Ortiz, R., 2016. Landrace  
 376 Germplasm for Improving Yield and Abiotic Stress Adaptation. *Trends Plant Sci.* 21, 31–42.  
 377 <https://doi.org/10.1016/j.tplants.2015.10.012>

378 Emebiri, L.C., 2013. QTL dissection of the loss of green colour during post-anthesis grain maturation  
 379 in two-rowed barley. *Theor. Appl. Genet.* 126, 1873–84. [https://doi.org/10.1007/s00122-013-](https://doi.org/10.1007/s00122-013-2102-0)  
 380 2102-0

381 Evans, L.T., 1997. Adapting and improving crops: the endless task. *Philos. Trans. R. Soc. B Biol. Sci.*  
 382 352, 901–906. <https://doi.org/10.1098/rstb.1997.0069>

383 Falster, D.S., Westoby, M., 2003. Leaf size and angle vary widely across species: What consequences  
 384 for light interception? *New Phytol.* 158, 509–525. [https://doi.org/10.1046/j.1469-](https://doi.org/10.1046/j.1469-8137.2003.00765.x)  
 385 8137.2003.00765.x

386 Farquhar, G.D., von Caemmerer S, Berry, J.A., 2001. Models of photosynthesis. *Plant Physiol.* 125,  
 387 42–45. <https://doi.org/10.1104/pp.125.1.42>

388 Fischer, R.A., Edmeades, G.O., 2010. Breeding and Cereal Yield Progress. *Crop Sci.*  
 389 <https://doi.org/10.2135/cropsci2009.10.0564>

390 Flood, P.J., Harbinson, J., Aarts, M.G.M., 2011. Natural genetic variation in plant photosynthesis.  
 391 *Trends Plant Sci.* <https://doi.org/10.1016/j.tplants.2011.02.005>

392 Gaju, O., DeSilva, J., Carvalho, P., Hawkesford, M.J., Griffiths, S., Greenland, A., Foulkes, M.J., 2016.  
 393 Leaf photosynthesis and associations with grain yield, biomass and nitrogen-use efficiency in  
 394 landraces, synthetic-derived lines and cultivars in wheat. *F. Crop. Res.* 193, 1–15.  
 395 <https://doi.org/10.1016/j.fcr.2016.04.018>

396 Giunta, F., Motzo, R., Deidda, M., 2002. SPAD readings and associated leaf traits in durum wheat,  
 397 barley and triticale cultivars. *Euphytica* 125, 197–205.

398 <https://doi.org/10.1023/a:1015878719389>

399 Goyne, P.J., Milroy, S.P., Lilley, J.M., Hare J M, 1993. Radiation Interception, Radiation Use Efficiency  
400 and Growth of Barley Cultivars. *Aust. J. Agric. Res.* 44, 1351–1366.

401 HGCA (The Scottish Executive), 2006. The barley growth guide.

402 Horton, P., Murchie, E.H., Chen, Y., Hubbart, S., Peng, S., Horton, P., 1999. Interactions between  
403 Senescence and Leaf Orientation Determine in Situ Patterns of Photosynthesis and  
404 Photoinhibition in Field-Grown Rice1. *Plant Physiol.* <https://doi.org/10.1104/pp.119.2.553>

405 Jones, H., Civiáň, P., Cockram, J., Leigh, F.J., Smith, L.M., Jones, M.K., Charles, M.P., Molina-Cano, J.-L.,  
406 Powell, W., Jones, G., Brown, T.A., 2011. Evolutionary history of barley cultivation in Europe  
407 revealed by genetic analysis of extant landraces. *BMC Evol. Biol.* 11, 320.  
408 <https://doi.org/10.1186/1471-2148-11-320>

409 Kemanian, A.R., Stöckle, C.O., Huggins, D.R., 2004. Variability of Barley Radiation-Use Efficiency. *Crop*  
410 *Sci.* <https://doi.org/10.2135/cropsci2004.1662>

411 Kruk, B., Insausti, P., Razul, A., Benech-Arnold, R., 2006. Light and thermal environments as modified  
412 by a wheat crop: effects on weed seed germination. *J. Appl. Ecol.* 43, 227–236.  
413 <https://doi.org/10.1111/j.1365-2664.2006.01140.x>

414 Ku, L.X., Zhao, W.M., Zhang, J., Wu, L.C., Wang, C.L., Wang, P.A., Zhang, W.Q., Chen, Y.H., 2010.  
415 Quantitative trait loci mapping of leaf angle and leaf orientation value in maize (*Zea mays* L.).  
416 *Theor. Appl. Genet.* 121, 951–959. <https://doi.org/10.1007/s00122-010-1364-z>

417 Kumagai, E., Hamaoka, N., Araki, T., Ueno, O., 2014. Dorsoventral asymmetry of photosynthesis and  
418 photoinhibition in flag leaves of two rice cultivars that differ in nitrogen response and leaf  
419 angle. *Physiol. Plant.* <https://doi.org/10.1111/ppl.12145>

420 Long, S.P., Marshall-Colon, A., Zhu, X.-G., 2015. Meeting the Global Food Demand of the Future by

421 Engineering Crop Photosynthesis and Yield Potential. *Cell* 161, 56–66.

422 <https://doi.org/10.1016/j.cell.2015.03.019>

423 Long, S.P., Zhu, X.-G., Naidu, S.L., Ort, D.R., 2006. Can improvement in photosynthesis increase crop

424 yields? *Plant. Cell Environ.* 29, 315–330. <https://doi.org/10.1111/j.1365-3040.2005.01493.x>

425 Lynch, J.P., Doyle, D., McAuley, S., McHardy, F., Danneels, Q., Black, L.C., White, E.M., Spink, J., 2017.

426 The impact of variation in grain number and individual grain weight on winter wheat yield in

427 the high yield potential environment of Ireland. *Eur. J. Agron.* 87, 40–49.

428 <https://doi.org/https://doi.org/10.1016/j.eja.2017.05.001>

429 Mackay, I., Horwell, A., Garner, J., White, J., McKee, J., Philpott, H., 2011. Reanalyses of the historical

430 series of UK variety trials to quantify the contributions of genetic and environmental factors to

431 trends and variability in yield over time. *Theor. Appl. Genet.* 122, 225–238.

432 <https://doi.org/10.1007/s00122-010-1438-y>

433 Madani, A., Rad, A.S., Pazoki, A., Nourmohammadi, G., Zarghami, R., 2010. Wheat (*Triticum aestivum*

434 L.) grain filling and dry matter partitioning responses to source:sink modifications under

435 postanthesis water and nitrogen deficiency. *Acta Sci. - Agron.* 32, 145–151.

436 <https://doi.org/10.4025/actasciagron.v32i1.6273>

437 Meseka, S., Fakorede, M., Ajala, S., Badu-Apraku, B., Menkir, A., 2013. Introgression of Alleles from

438 Maize Landraces to Improve Drought Tolerance in an Adapted Germplasm. *J. Crop Improv.* 27,

439 96–112. <https://doi.org/10.1080/15427528.2012.729259>

440 Meseka, S., Menkir, A., Obeng-Antwi, K., 2015. Exploitation of beneficial alleles from maize (*Zea*

441 mays L.) landraces to enhance performance of an elite variety in water stress environments.

442 *Euphytica* 201, 149–160. <https://doi.org/10.1007/s10681-014-1214-1>

443 Monteith, J.L., Moss, C.J., 1977. Climate and the Efficiency of Crop Production in Britain [and

444 Discussion]. *Philos. Trans. R. Soc. B Biol. Sci.* <https://doi.org/10.1098/rstb.1977.0140>

445 Mukula, J., Rantanen, O., 1837. Climatic risks to the yield and quality of field crops in Finland: I. Basic  
446 facts about Finnish field crops production. *Ann. Agric. Fenn.* 26, 1–18.

447 Murchie, E.H., Pinto, M., Horton, P., 2009. Agriculture and the new challenges for photosynthesis  
448 research. *New Phytol.* 181, 532–552. <https://doi.org/10.1111/j.1469-8137.2008.02705.x>

449 Muurinen, S., Peltonen-Sainio, P., 2006. Radiation-use efficiency of modern and old spring cereal  
450 cultivars and its response to nitrogen in northern growing conditions. *F. Crop. Res.* 96, 363–  
451 373. <https://doi.org/10.1016/j.fcr.2005.08.009>

452 Nan Su San, Ootsuki, Y., Adachi, S., Yamamoto, T., Ueda, T., Tanabata, T., Motobayashi, T., Ookawa,  
453 T., Hirasawa, T., 2018. A near-isogenic rice line carrying a QTL for larger leaf inclination angle  
454 yields heavier biomass and grain. *F. Crop. Res.* 219, 131–138.  
455 <https://doi.org/10.1016/j.fcr.2018.01.025>

456 Nunes-Nesi, A., Nascimento, V. de L., de Oliveira Silva, F.M., Zsögön, A., Araújo, W.L., Sulpice, R.,  
457 2016. Natural genetic variation for morphological and molecular determinants of plant growth  
458 and yield. *J. Exp. Bot.* 67, 2989–3001. <https://doi.org/10.1093/jxb/erw124>

459 Ort, D.R., Merchant, S.S., Alric, J., Barkan, A., Blankenship, R.E., Bock, R., Croce, R., Hanson, M.R.,  
460 Hibberd, J.M., Long, S.P., Moore, T.A., Moroney, J., Niyogi, K.K., Parry, M.A.J., Peralta-Yahya,  
461 P.P., Prince, R.C., Redding, K.E., Spalding, M.H., van Wijk, K.J., Vermaas, W.F.J., von Caemmerer,  
462 S., Weber, A.P.M., Yeates, T.O., Yuan, J.S., Zhu, X.G., 2015. Redesigning photosynthesis to  
463 sustainably meet global food and bioenergy demand. *Proc. Natl. Acad. Sci.* 112, 8529–8536.  
464 <https://doi.org/10.1073/pnas.1424031112>

465 Ort, D.R., Zhu, X., Melis, A., 2011. Optimizing Antenna Size to Maximize Photosynthetic Efficiency.  
466 *Plant Physiol.* 155, 79–85. <https://doi.org/10.1104/pp.110.165886>

467 Parry, M.A.J., Reynolds, M., Salvucci, M.E., Raines, C., Andralojc, P.J., Zhu, X.G., Price, G.D., Condon,  
 468 A.G., Furbank, R.T., 2011. Raising yield potential of wheat. II. Increasing photosynthetic capacity  
 469 and efficiency. *J. Exp. Bot.* 62, 453–467. <https://doi.org/10.1093/jxb/erq304>

470 Peltonen-Sainio, P., 1997. Leaf Area Duration of Oat at High Latitudes. *J. Agron. Crop Sci.* 178, 149–  
 471 155. <https://doi.org/10.1111/j.1439-037X.1997.tb00483.x>

472 Pendleton, J.W., Smith, G.E., Winter, S.R., Johnston, T.J., 1968. Field Investigations of the  
 473 Relationships of Leaf Angle in Corn (*Zea mays* L.) to Grain Yield and Apparent Photosynthesis1.  
 474 *Agron. J.* 60, 422. <https://doi.org/10.2134/agronj1968.00021962006000040027x>

475 Peng, S., Khush, G.S., Virk, P., Tang, Q., Zou, Y., 2008. Progress in ideotype breeding to increase rice  
 476 yield potential. *F. Crop. Res.* <https://doi.org/10.1016/j.fcr.2008.04.001>

477 Raines, C.A., 2011. Increasing photosynthetic carbon assimilation in C3 plants to improve crop yield:  
 478 current and future strategies. *Plant Physiol.* 155, 36–42.  
 479 <https://doi.org/10.1104/pp.110.168559>

480 Ray, D.K., Ramankutty, N., Mueller, N.D., West, P.C., Foley, J. a, 2012. Recent patterns of crop yield  
 481 growth and stagnation. *Nat. Commun.* 3, 1293. <https://doi.org/10.1038/ncomms2296>

482 Reynolds, M., Bonnett, D., Chapman, S.C., Furbank, R.T., Manès, Y., Mather, D.E., Parry, M.A.J., 2011.  
 483 Raising yield potential of wheat. I. Overview of a consortium approach and breeding strategies.  
 484 *J. Exp. Bot.* 62, 439–452. <https://doi.org/10.1093/jxb/erq311>

485 Reynolds, M., Foulkes, M.J., Slafer, G.A., Berry, P., Parry, M.A.J., Snape, J.W., Angus, W.J., 2009.  
 486 Raising yield potential in wheat. *J. Exp. Bot.* 60, 1899–1918.  
 487 <https://doi.org/10.1093/jxb/erp016>

488 Richards, R.A., 2000. Selectable traits to increase crop photosynthesis and yield of grain crops. *J. Exp.*  
 489 *Bot.* 51, 447–458. [https://doi.org/10.1093/jexbot/51.suppl\\_1.447](https://doi.org/10.1093/jexbot/51.suppl_1.447)

490 Rodriguez, M., Rau, D., Papa, R., Attene, G., 2008. Genotype by environment interactions in barley  
 491 (*Hordeum vulgare* L.): different responses of landraces, recombinant inbred lines and varieties  
 492 to Mediterranean environment. *Euphytica* 163, 231–247. [https://doi.org/10.1007/s10681-007-](https://doi.org/10.1007/s10681-007-9635-8)  
 493 9635-8

494 Ryel, R.J., Beyschlag, W., Caldwell, M.M., 1993. Foliage orientation and carbon gain in two tussock  
 495 grasses as assessed with a new whole-plant gas-exchange model. *Funct. Ecol.*  
 496 <https://doi.org/10.2307/2389874>

497 Serrago, R.A., Alzueta, I., Savin, R., Slafer, G.A., 2013. Understanding grain yield responses to source-  
 498 sink ratios during grain filling in wheat and barley under contrasting environments. *F. Crop. Res.*  
 499 150, 42–51. <https://doi.org/10.1016/j.fcr.2013.05.016>

500 Sim, L.C., Froud-Williams, R.J., Gooding, . J, 2007. The influence of winter oilseed rape (*Brassica*  
 501 *napus* ssp. *oleifera* var. *biennis*) canopy size on grass weed growth and grass weed seed return.  
 502 *J. Agric. Sci.* 145, 313–327. <https://doi.org/10.1017/S0021859606006721>

503 Taylaran, R.D., Adachi, S., Ookawa, T., Usuda, H., Hirasawa, T., 2011. Hydraulic conductance as well  
 504 as nitrogen accumulation plays a role in the higher rate of leaf photosynthesis of the most  
 505 productive variety of rice in Japan. *J. Exp. Bot.* 62, 4067–4077.  
 506 <https://doi.org/10.1093/jxb/err126>

507 The Scottish Government, 2018. Cereal and oilseed rape harvest: 2018 final estimates.

508 Turner, A., Beales, J., Faure, S., Dunford, R.P., Laurie, D.A., 2005. The pseudo-response regulator Ppd-  
 509 H1 provides adaptation to photoperiod in barley. *Science* (80-. ). 310, 1031–1034.  
 510 <https://doi.org/10.1126/science.1117619>

511 Valladares, F., Pugnaire, F.I., 1999. Tradeoffs between irradiance capture and avoidance in semi-arid  
 512 environments assessed with a crown architecture model. *Ann. Bot.*

513        <https://doi.org/10.1006/anbo.1998.0843>

514    Villa, T.C.C., Maxted, N., Scholten, M., Ford-Lloyd, B., 2005. Defining and Identifying Crop Landraces.  
515        Plant Genet. Resour. 3, 373–384. <https://doi.org/10.1079/PGR200591>

516    Werner, C., Ryel, R.J., Correia, O., Beyschlag, W., 2001. Structural and functional variability within the  
517        canopy and its relevance for carbon gain and stress avoidance. Acta Oecologica.  
518        [https://doi.org/10.1016/S1146-609X\(01\)01106-7](https://doi.org/10.1016/S1146-609X(01)01106-7)

519    Xu, K., Mackill, D.J., 1996. A major locus for submergence tolerance mapped on rice chromosome 9.  
520        Mol. Breed. 2, 19–224. <https://doi.org/10.1007/BF00564199>

521    Yahiaoui, S., Cuesta-Marcos, A., Gracia, M.P., Medina, B., Lasa, J.M., Casas, A.M., Ciudad, F.J.,  
522        Montoya, J.L., Moralejo, M., Molina-Cano, J.L., Igartua, E., 2014. Spanish barley landraces  
523        outperform modern cultivars at low-productivity sites. Plant Breed. 133, 218–226.  
524        <https://doi.org/10.1111/pbr.12148>

525    Yin, X., Struik, P.C., 2015. Constraints to the potential efficiency of converting solar radiation into  
526        phytoenergy in annual crops: from leaf biochemistry to canopy physiology and crop ecology. J.  
527        Exp. Bot. 66, 6535–6549. <https://doi.org/10.1093/jxb/erv371>

528    Zadoks, J.C., Chang, T.T., Konzak, C.F., 1974. A decimal code for the growth stages of cereals. Weed  
529        Res. 14, 415–421. <https://doi.org/10.1111/j.1365-3180.1974.tb01084.x>

530    Zhang, L., Richards, R.A., Condon, A.G., Liu, D.C., Rebetzke, G.J., 2015. Recurrent selection for wider  
531        seedling leaves increases early biomass and leaf area in wheat (*Triticum aestivum* L.). J. Exp.  
532        Bot. 66, 1215–1226. <https://doi.org/10.1093/jxb/eru468>

533    Zheng, H.J., Wu, a. Z., Zheng, C.C., Wang, Y.F., Cai, R., Shen, X.F., Xu, R.R., Liu, P., Kong, L.J., Dong,  
534        S.T., 2009. QTL mapping of maize ( *Zea mays* ) stay-green traits and their relationship to yield.  
535        Plant Breed. 128, 54–62. <https://doi.org/10.1111/j.1439-0523.2008.01529.x>



536 Zhu, X.-G., Long, S.P., Ort, D.R., 2010. Improving photosynthetic efficiency for greater yield. *Annu.*  
537 *Rev. Plant Biol.* 61, 235–261. <https://doi.org/10.1146/annurev-arplant-042809-112206>

538